Appl. No. 10/056,212 Amdt. dated October 1, 2003 Reply to Office action of July 2, 2003

Amendments to the Claims

comprising:

This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Currently Amended) A positive working imageable composition,

an ink-receptive hydroxyfunctional resin comprising a covalently bound radiation sensitive group capable of increasing the solubility of said imageable composition in an alkaline developer upon exposure to radiation; and

an isocyanate crosslinking agent.

2. (Cancelled)

 (Currently Amended) The imageable composition of claim 1, wherein said covalently bound radiation sensitive group is derived from an sensitive to ultraviolet radiation sensitive compound.

4. (Currently Amended) The imageable composition of claim 3, wherein said ultraviolet radiation sensitive compound group is selected from the group consisting of compounds represented by moieties of the formula:

wherein

Appl. No. 10/056,212 Amdr. dated October 1, 2003 Reply to Office action of July 2, 2003

each of R¹, R², R³, R⁴ and R⁵ is independently selected from the group consisting of: hydrogen, linear, branched or cyclic alkyl of 1 to 22 carbon atoms, aralkyl of 6 to 22 carbon atoms, aryl, alkaryl, alkoxy of 1 to 22 carbon atoms, haloalkyl, halogen, acyl, ester and cyano, and

wherein a is halogen.

5. (Currently Amended) The imageable composition of claim 4, wherein said altraviolet radiation sensitive compound group is selected from the group consisting of compounds represented by moieties of the formula:

wherein

u is chlorine.

- 6. (Cancelled)
- 7. (Currently Amended) The imageable composition of claim 6 1, wherein said polyfunctional phenolic resin free of radiation sensitive groups is hydroxyfunctional resin further comprises a resin moiety derived from a polyfunctional resin selected from the group consisting of: polyfunctional phenolic resin, a novolak resin, a pyrogallol/acetone resin, polyvinyl phenol polymer, vinyl phenol/hydrocarbyl acrylate copolymer, a resole resin, an acrylic resin, a polyester resin, a polyurethane resin, a polyol and a mixture thereof.
- 8. (Currently Amended) The imageable composition of claim 7, wherein said polyfunctional phenolic resin free of radiation sensitive groups polyfunctional resin is selected from the group consisting of a phenol novolak resin, a cresol novolak resin, a phenol/cresol novolak resin, a resole resin and or a mixture thereof.

9-12. (Cancelled)

13. (Currently Amended) The imageable composition f claim 11. wherein said isocyanate crosslinking agent is selected from the group consisting of: isophorone



Appl. No. 10/056,212 Amdt dated October 1, 2003 Reply to Office action of July 2, 2003

f d

diisocyanate, methylene-bis-phenyl diisocyanate, toluene diisocyanate, hexamethylene diisocyanate, tetramethylxylylene diisocyanate, dimers thereof, adducts thereof with diols, adducts thereof with triols, adducts thereof with polyols, adducts thereof with polyesters, adducts thereof with acrylic resins, adducts thereof with polyurethane polyols, adducts thereof with an isocyanate blocking agent and mixtures thereof.

- 14. (Currently Amended) The imageable composition of claim 13, wherein said isocyanate <u>crosslinking agent further comprises a</u> blocking agent is selected from the group consisting of: a phenol, an oxime, a lactam and a pyrazole.
- 15. (Currently Amended) The imageable composition of claim 14, wherein said isocyanate blocking agent is selected from the group consisting of phenol, methyl ethyl ketone oxime, 2-pytrolidone, 2-piperidone, caprolactam and or 3,5-dimethylpyrazole.
 - 16. (Original) The imageable composition of claim 1, further comprising: a colorant; and an acid generator.
- 17. (Original) The composition of claim 16, wherein said colorant is selected from the group consisting: of a colorant dye, a colorant pigment and a combination thereof.
- 18. (Currently Amended) The composition of claim 17, wherein said colorant dye is selected from the group consisting of: crystal violet, crystal violet lactone, basonyl blue, victoria pure blue BO, victoria blue B, blue colorant dye victoria blue FBR represented by the formula:




Appl, No. 10/056,212 Amdt, dated October 1, 2003 Reply to Office action of July 2, 2003

19. (Cancelled)

20. (Currently Amended) The imageable composition of claim 19 16, wherein said acid generator is a light sensitive triazine compound represented by of the formula:

wherein

Z selected from the group consisting of: hydrogen, linear, branched or cyclic alkyl of 1 to 22 carbon atoms, aralkyl of 6 to 22 carbon atoms, aryl, alkaryl, alkoxy of 1 to 22 carbon atoms, haloalkyl, halogen, acyl, ester, cyano, a group represented by moiety of the formula:



Page 5 of 12

Appl. No. 10/056,212 Arndt, dated October 1, 2003 Reply to Office action of July 2, 2003

wherein

each of R⁶, R⁷, R⁸ and R⁹ is independently selected from the group consisting of: hydrogen, linear, branched or cyclic alkyl of 1 to 22 carbon atoms, aralkyl of 6 to 22 carbon atoms, aryl, alkaryl, alkoxy of 1 to 22 carbon atoms, haloalkyl, halogen, acyl, ester and cyano, wherein or R⁶ and R⁷ or end/or R⁸ and R⁹ together with carbon atoms to which they are attached to can form a cycloaliphatic, benzo or a substituted benzo ring;

wherein R¹⁰ is selected from the group consisting of: linear, branched or cyclic alkyl of 1 to 22 carbon atoms, aralkyl of 6 to 22 carbon atoms, aryl, alkaryl, haloalkyl, acyl, ester and cyano; and wherein Y is a heterostem selected from the group consisting of:

oxygen and or sulfur.

21. (Currently Amended) The imageable composition of claim 20, wherein light sensitive triazine compound is represented by of the formula:

wherein


each of R⁶, R⁷, R⁸ and R⁹ is independently selected from the group consisting of:
hydrogen, linear, branched or cyclic alkyl of 1 to 22 carbon atoms, aralkyl
of 6 to 22 carbon atoms, aryl, alkaryl, alkoxy of 1 to 22 carbon atoms,
haloalkyl, halogen, acyl, ester and cyano, wherein or R⁶ and R⁷ or and/or
R⁸ and R⁹ together with carbon atoms to which they are attached to
ean form a cycloaliphatic, benzo or a substituted benzo ring;


wherein R¹⁰ is selected from the group consisting of: linear, branched or cyclic alkyl of 1 to 22 carbon atoms, aralkyl of 6 to 22 carbon atoms, aryl, alkaryl, haloalkyl, acyl, ester and cyano; and


Appl. No. 10/056,212 Amdt. dated October 1, 2003 Reply to Office action of July 2, 2003

wherein Y is a heterestem selected from the group consisting of xygen and or sulfur.

The imageable composition of claim 21, wherein said light 22. (Original) sensitive triazine compound is selected from the group consisting of:







and a mixture thereof.

(Currently Amended) The imageable composition of claim 19 16, wherein 23. said onium salt acid generator is selected from the group consisting of: an iodonium salt, a sulfonium salt, a hydrocarbyloxysulfonium salt, a hydrocarbyloxyammonium salt, an aryl diazonium salt and a combination thereof.

24-33. (Cancelled)

- (Currently Amended) An imageable element comprising: 34.
- a substrate comprising a hydrophilic surface; and
- a positive working imageable composition coated on a surface of said substrate, said composition comprising:
 - a an ink-receptive hydroxyfunctional resin comprising a covalently bound radiation sensitive group capable of increasing the solubility of said imageable composition in an alkaline developer upon exposure to radiation; and

an isocyanate crosslinking agent.

- 35. (Cancelled)
- (Original) The imageable element of claim 34, wherein said positive 36. working imageable composition further comprises a colorant and an acid generator.

Appl. No. 10/056,212 Amdt. dated October 1, 2003 Reply to Office action of July 2, 2003

37. (Currently Amended) A method of producing an imaged element comprising the steps of:

providing an imageable element comprising a substrate <u>comprising a</u>

<u>hydrophilic surface</u> and a positive working imageable composition coated on a <u>the</u> surface of said substrate, <u>wherein</u> said <u>imageable</u> composition comprising <u>comprises</u>:

an ink-receptive hydroxyfunctional resin comprising a covalently bound radiation sensitive group capable of increasing the solubility of said imageable composition in an alkaline developer upon exposure to radiation; and

an isocyanate crosslinking agent;

heating said imageable element at a temperature and length of time sufficient to produce a crosslinked imageable element;

imagewise exposing said crosslinked imageable element to radiation to produce an imagewise exposed element having exposed and unexposed regions; and

removing the exposed regions of said imageable composition and thereby to produce said imaged element.

- 38. (Original) The method of claim 37, wherein said positive working imageable composition further comprises a colorant and an acid generator.
- 39. (Original) The method of claim 37, wherein said exposing step is carried out using ultraviolet radiation.

40-49. (Cancelled)